

THREEFOLDS OF $\mathbb{P}^5$ WITH ONE APPARENT QUADRUPLER POINT

PIETRO DE POI

Dipartimento di Scienze Matematiche
 Università degli Studi di Trieste
 Via Valerio, 12/b
 I-34127 Trieste
 Italy
 depoi@mathsun1.univ.trieste.it

ABSTRACT. In this article we classify all the smooth threefolds of $\mathbb{P}^5$ with an apparent quadruple point provided that the family of its 4-secant lines is an irreducible (first order) congruence. This is sufficient to conclude the classification of all the smooth codimension two varieties of $\mathbb{P}^n$ with one apparent $(n-1)$ -point and with irreducible family of $(n-1)$ -secant lines.

INTRODUCTION

A congruence of lines in $\mathbb{P}^n$ is a family of lines of dimension $n-1$, and its order is the number of lines passing through a general point of $\mathbb{P}^n$. A codimension two subvariety of $\mathbb{P}^n$ is said to have q apparent $(n-1)$ -tuple points if its general projection from a point to a hyperplane has q $(n-1)$ -tuple points as singularities.

In this work we prove that the degree of these varieties is bounded by $(n-1)^2$ when they have only one apparent $(n-1)$ -tuple point (*i.e.* $q=1$), and we observe that this implies that they cannot be complete intersections. By a result of A. Holme and M. Schneider, [HS85], this implies also that in order to classify the smooth ones, we can stop up to dimension three.

In this paper we also give a partial result towards this classification, *i.e.* we restrict ourselves to the case in which the family of the $(n-1)$ -secant lines is in fact an irreducible first order congruence.

This article is structured as follows: after giving, in Section 1, the basic definitions, we obtain some general results about first order congruences in $\mathbb{P}^n$. In particular, after giving the central definition of fundamental d -loci, we show how to obtain the degree bound for the fundamental $(n-2)$ -locus of a congruence.

In Section 2, we give two general examples of congruences in $\mathbb{P}^n$: the first one is that of linear congruences, *i.e.* the congruences which come out from general linear sections of the Grassmannians; the second one is that of congruences given by the $(n-1)$ -secant lines of the varieties given by the degeneracy locus of a general map $\phi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}, \mathcal{O}_{\mathbb{P}^n}^{\oplus n}(1))$. Of these two examples we calculate locally free resolutions of the ideal sheaves of their focal locus and of the congruences themselves. We study these examples because they give us all the congruences of our classification but one (*i.e.* case (4b) of Theorem 0.1).

In Section 3 we prove two multiple point formulae: the quadruple point formula for a smooth threefold of $\mathbb{P}^5$ and the formula which gives the number of 4-secant

1991 *Mathematics Subject Classification.* Primary 14J30, 14M15, 14N05, Secondary 51N35.

Key words and phrases. Algebraic Geometry, threefolds, Focal loci, Congruences.

The author has a Post-Doc grant at the Trieste University and is a member of GNSAGA-CNR.

lines to a smooth surface of $\mathbb{P}^4$ passing through a general point of the surface itself. Strangely enough, the quadruple point formula for the threefolds—which is actually an application of S. Kleiman’s multiple point formulae of maps (see [Kle81])—it seems to have been unknown, at least in modern times (see for example [BSS95]). With this formula, E. Mezzetti has been able to exclude the only degree 12 smooth threefold of $\mathbb{P}^5$ for which the existence was uncertain (see [Ede94]). I must say that after the proof of this formula, I realized that the same was obtained by S. Kwak in [Kwa01], but with other methods, *i.e.* through the monoidal construction.

The irreducible congruences of order one of $\mathbb{P}^5$ which are given by the 4-secant lines of a smooth threefold are classified in Section 4. By the results of Section 2 and what we said above, we obtain the following complete list, where d is the degree of the smooth codimension two variety X , π its sectional genus and, if $\dim X = 3$, S is its general hyperplane section; finally, H and K are the hyperplane and canonical divisor classes, respectively (and X_i is referred to the classification given in [DP95]):

Theorem 0.1. *The smooth codimension two (irreducible) subvarieties of $\mathbb{P}^n$ for which the family of their $(n-1)$ -secant lines is an irreducible first order congruence are*

1. *for $n = 2$ a point, and the congruence is a pencil of lines;*
2. *for $n = 3$ the twisted cubic, and the congruence is the Veronese surface and has bidegree $(1, 3)$ (for more details, see Subsection 2.2);*
3. *for $n = 4$ we have the following possibilities:*
 - (a) *a (projected) Veronese surface, which is rational, with $d = 4$, $\pi = 0$; in this case we have a linear congruence, which has bidegree $(1, 2)$ (see Subsection 2.1);*
 - (b) *a Bordiga surface, which is rational, with $d = 6$, $\pi = 3$; the congruence is smooth and has bidegree $(1, 8)$ (see Subsection 2.2);*
4. *for $n = 5$ we have the following possibilities:*
 - (a) *the Palatini scroll, which is rational with $d = 7$, $\pi = 4$, $\chi(\mathcal{O}_S) = \chi(\mathcal{O}_X) = 1$ (case X_6); in this case we have a linear congruence, which has 3-degree $(1, 3, 2)$ (see Subsection 2.1);*
 - (b) *a non rational scroll, $\mathbb{P}^1$ -bundle over a minimal K3 surface of $\mathbb{P}^8$ via $|K + H|$ (case X_{11}); $d = 9$, $\pi = 8$, $\chi(\mathcal{O}_S) = \chi(\mathcal{O}_X) = 2$; the congruence, has 3-degree $(1, 7, 13)$;*
 - (c) *a log-general type rational threefold, linked with a $(4, 4)$ complete intersection to a Bordiga threefold (case X_{15}); $d = 10$, $\pi = 11$, $\chi(\mathcal{O}_S) = 5$, $\chi(\mathcal{O}_X) = 1$; the congruence is smooth (see Subsection 2.2) and has 3-degree $(1, 15, 20)$.*

Vice versa the $(n-1)$ -secant lines of any of the above varieties generate a first order congruence.

We conjecture that the first order congruences whose pure fundamental locus F (see Definition 1.1 below) is a smooth variety of codimension two are the ones obtained in Theorem 0.1 (*i.e.* the families of $(n-1)$ -secant lines of the varieties of the theorem).

Acknowledgments. I would like to thank Emilia Mezzetti, Ragni Piene, Kristian Ranestad, Fyodor L. Zak and Jon Eivind Vatne for interesting comments and remarks.

I also thank the referee for the careful reading and important suggestions and remarks.

1. NOTATIONS, DEFINITIONS AND GENERAL RESULTS

We will work with schemes and varieties over the complex field $\mathbb{C}$. By *variety* we mean a reduced and irreducible algebraic $\mathbb{C}$ -scheme. More information about general results and references about families of lines, focal diagrams and congruences can be found in [De 01] or [De 99]. Besides, we refer to [GH78] for notations about Schubert cycles and to [Ful84] for the definitions and results of intersection theory. Here we recall that a *congruence of lines* of $\mathbb{P}^n$ is a flat family (Λ, B, p) of lines of $\mathbb{P}^n$ obtained by the desingularization of a subvariety B' of dimension $n - 1$ of the Grassmannian $\mathbb{G}(1, n)$ of lines of $\mathbb{P}^n$. p is the restriction of the projection $p_1 : B \times \mathbb{P}^n \rightarrow B$ to Λ , while we will denote the restriction of $p_2 : B \times \mathbb{P}^n \rightarrow \mathbb{P}^n$ by f . $\Lambda_b := p^{-1}(b)$, ($b \in B$) will be an element of the family and $f(\Lambda_b) =: \Lambda(b)$ is a line of $\mathbb{P}^n$. We can summarise all these notations in the following two diagrams: the first one defines the family

$$\begin{array}{ccccc} \Lambda := \psi^*(\mathcal{H}_{1,n}) & \xrightarrow{\psi^*} & \mathcal{H}_{1,n} & \xrightarrow{p_2} & \mathbb{P}^n \\ p \downarrow & & p_1 \downarrow & & \\ B & \xrightarrow{\psi} & B' \subset \mathbb{G}(1, n), & & \end{array}$$

where $\mathcal{H}_{1,n} \subset \mathbb{G}(1, n) \times \mathbb{P}^n$ is the incidence variety and ψ is the desingularization map, and the second one explains the notation for the elements of the family

$$\begin{array}{ccc} \Lambda_b \subset \Lambda & \xrightarrow{f := \psi^* p_2} & \mathbb{P}^n \supset \Lambda(b) := f(\Lambda_b) \\ p \downarrow & & \\ b \in B. & & \end{array}$$

A point $y \in \mathbb{P}^n$ is called *fundamental* if its fibre $f^{-1}(y)$ has dimension greater than the dimension of the general one. The *fundamental locus* is the set of the fundamental points. The *subscheme of the foci of the first order*, or, simply, the *focal subscheme* $V \subset \Lambda$ is the scheme of the ramification points of f . The *locus of the first order foci*, or, simply, the *focal locus*, $\Phi := f(V) \subset \mathbb{P}^n$, is the set of the branch points of f . In this article, as we did in [De 01], we will endow this locus with the scheme structure given by considering it as the scheme-theoretic image of V under f (see, for example, [Har77]).

To a congruence is associated a *sequence of degrees* or $(\nu + 1)$ -*degree* $(a_0, \dots, a_\nu)$ if we write

$$[B] = \sum_{i=0}^{\nu} a_i \sigma_{n-1-i,i}$$

—where we put $\nu := \lfloor \frac{n-1}{2} \rfloor$ —as a linear combination of Schubert cycles of the Grassmannian (recall that $\sigma_{n-1-i,i} := [\{\ell \in \mathbb{G}(1, n) \mid \ell \cap \mathbb{P}^i \neq \emptyset; \ell, \mathbb{P}^i \subset \mathbb{P}^{n-i}\}]$). In particular, the *order* a_0 is the number of lines of B passing through a general point of $\mathbb{P}^n$. The fundamental locus is contained in the focal locus and the two loci coincide in the case of a first order congruence, *i.e.* through a focal point there will pass infinitely many lines of the congruence. An important result—independent of order and class—is the following (see also [De 01]):

Proposition 1.1. *On every line $\Lambda_b \subset \Lambda$ of the family, the focal subscheme V either coincides with the whole Λ_b —in which case $\Lambda(b)$ is called focal line—or is a zero dimensional subscheme of Λ_b of length $n - 1$. Moreover, in the latter case, if Λ is a first order congruence, $\Phi \cap \Lambda_b$ has length $n - 1$.*

Proof. Let $\lambda : \mathcal{T}_{(B \times \mathbb{P}^n / \mathbb{P}^n)|_\Lambda} \rightarrow \mathcal{N}_{\Lambda/B \times \mathbb{P}^n}$ be the *global characteristic map* for the family Λ (see [CS89]). From the *focal diagram* (diagram (3) of [CS89]) one gets

that the subscheme of the foci of the first order V is the degeneracy locus of λ . If we restrict the map λ to a fibre $\Lambda_b \cong \Lambda(b)$, we obtain the *characteristic map of the family relative to b* :

$$\begin{array}{ccc} \lambda(b) : T_{B,b} \otimes \mathcal{O}_{\Lambda(b)} & \longrightarrow & \mathcal{N}_{\Lambda(b)/\mathbb{P}^n} \\ \cong \downarrow & & \cong \downarrow \\ \mathcal{O}_{\Lambda(b)}^{n-1} & \longrightarrow & \mathcal{O}_{\Lambda(b)}(1)^{n-1}. \end{array}$$

From the preceding isomorphisms, the map $\lambda(b)$ can be seen as an $(n-1) \times (n-1)$ -matrix with linear entries on $\Lambda(b)$; so the focal locus on $\Lambda(b)$ is given by the vanishing of the determinant of this matrix, and our claim follows.

Concerning the first order congruences, we observe that a fundamental point P is a focal point for every line $\Lambda(b)$ which contains it, since the characteristic map relative to it, $\lambda(b)$, drops rank in P . $\square$

Remark. If Λ is a congruence of order different from one and the intersection $F \cap \Lambda(b)$ is proper, this can have length greater than $n-1$. This is due to the fact that $\Lambda(b)$ can contain points of Φ that are actually focal for other lines, *i.e.* points that are images of points contained in some $V \cap \Lambda_{b'}$, with $b \neq b'$.

From now on we will consider only first order congruences, and since in this case the focal and the fundamental loci coincide, we will talk only of fundamental loci (with the scheme structure given above for the focal locus). A central definition, introduced first in [De 01], is that of *fundamental d -locus*, *i.e.* the subscheme of the fundamental locus of pure dimension d , with $0 \leq d \leq n-2$, which is met by the general line of the congruence. Let us see how these schemes are constructed: the closed set

$$S_d := \{(\Lambda(b), P) \in \Lambda \mid \text{rk}(\text{d}f_{(\Lambda(b), P)}) \leq d\}$$

has a natural subscheme structure, which is defined by a Fitting ideal, *i.e.* the ideal generated by the $(d+1)$ -minors of $\text{d}f$ (or, by the Fitting lemma, see [Eis95], by the d -minors of λ), see [Kle77]; in particular, $S_{n-1} = V$. Let us define

$$D_{d+1} := \overline{S_{d+1} \setminus S_d}$$

with the scheme structure induced by S_{d+1} . Finally, we consider the scheme-theoretic image Φ_d of D_{d+1} in $\mathbb{P}^n$ under f . The component of Φ_d of pure dimension d (with the scheme structure induced by Φ_d) which is met by the general line of the congruence is the fundamental d -locus.

These subschemes of the fundamental locus are particularly important, since a first order congruence can be characterised as a component of the set of lines which meet the fundamental d -loci a certain number of times, see the Classification Theorem 3.2 of [De 01].

Definition 1.1. The union of the fundamental d -loci of F is called *pure fundamental locus*, or, in what follows, simply *fundamental locus* and it is denoted by F .

After this, we give the following

Theorem 1.2. *If Λ is a first order congruence such that the pure fundamental locus F is irreducible and coincides with the fundamental $(n-2)$ -locus, then*

$$\frac{n-1}{k} < m < (n-1)^2,$$

where $m := \deg(F)_{\text{red}}$ and k is the geometric multiplicity $(F)_{\text{red}}$ in F .

Proof. First of all, we have that $n-1 < km$ by degree reasons, since the congruence is given by lines which intersect F in a zero dimensional scheme of length $n-1$.

To prove the other bound, we need of course more work. Let B be our congruence, which has sequence of degrees $(1, a_1, \dots, a_\nu)$. If Π is a (fixed) general $(n-2)$ -plane, we denote by V_Π the scroll given by the lines of the congruence that meet Π . Then by the Schubert calculus one can show that V_Π is a hypersurface of $\mathbb{P}^n$ of degree $1 + a_1$.

Moreover, if ℓ is a line of B not contained in V_Π and P is a point of $V_\Pi \cap \ell$, then P is a focus for B , since at least two lines of the congruence pass through it.

Then, if Π' is another general $(n-2)$ -plane of $\mathbb{P}^n$, the complete intersection of the hypersurfaces V_Π and $V_{\Pi'}$ is a (reducible) $(n-2)$ -dimensional scheme Γ that contains the focal locus F and the $(n-2)$ -dimensional scroll Σ given by the lines of the congruence meeting Π and Π' , which has degree $1 + 2a_1 + a_2$ (actually, $1 + a_1$ in the case of $\mathbb{P}^3$; since this case can be treated analogously, we will suppose from now on that $n > 3$).

In fact, if a point P of $V_\Pi \cap V_{\Pi'}$ does not belong to the scroll Σ , it belongs to the fundamental locus. Indeed in this case

$$P \in \ell \cap \ell', \text{ where } \ell \in G_\Pi, \ell \in G_{\Pi'}, \text{ and } \ell \neq \ell'$$

—where G_Π and $G_{\Pi'}$ denote the subvarieties of the Grassmannian corresponding to the two scrolls V_Π and $V_{\Pi'}$. Since Λ is a first order congruence and P belongs to two of the lines of Λ , it belongs to infinitely many ones.

Reciprocally, if $P \in F$ is a general focal point, the set of the lines of B through P , χ_P , is a cone of dimension (at least) two, so its intersection with Π and Π' is not empty and therefore $P \in V_\Pi \cap V_{\Pi'}$.

The degree of the scroll follows from the Schubert calculus.

Claim. The following formulae hold:

$$\begin{aligned} (1) \quad & (n-1)h = 1 + a_1, \\ (2) \quad & (1 + a_1)^2 \geq h^2m + 1 + 2a_1 + a_2, \end{aligned}$$

where we denoted by h the algebraic multiplicity of $(F)_{\text{red}}$ on V_Π .

Let us start proving relation (1). If we take a line $\Lambda(b)$ of the congruence not contained in $F \cap V_\Pi$, then, intersecting $\Lambda(b)$ with V_Π , we obtain a zero dimensional scheme of length $1 + a_1$, since this is the degree of V_Π . $V_\Pi \cap \Lambda(b)$, set-theoretically, coincides with $(F)_{\text{red}} \cap \Lambda(b)$, and its intersection multiplicity is h , so the relation (1) is proved.

We recall that the degree of Γ is $(1 + a_1)^2$, and it contains F and the scroll Σ . Actually, the fundamental $(n-2)$ -locus has geometric multiplicity in Γ equal to h^2 . The proof of this fact is the following: the intersection multiplicity $i((F)_{\text{red}}, V_\Pi \cdot V_{\Pi'}, \mathbb{P}^n)$ of F in $V_\Pi \cdot V_{\Pi'}$ is equal to the geometric multiplicity of $(F)_{\text{red}}$ in Γ , but $i((F)_{\text{red}}, V_\Pi \cdot V_{\Pi'}, \mathbb{P}^n) = h^2$.

Finally, as we seen, the scroll Σ has degree $1 + 2a_1 + a_2$, so we get formula (2).

Now, if we substitute formula (1) in formula (2), we obtain

$$(3) \quad (n-1)^2h^2 - h^2m - 1 - 2a_1 - a_2 \geq 0,$$

and since $-1 - 2a_1 - a_2 < 0$, we deduce $m < (n-1)^2$. $\square$

A fundamental consequence of the preceding theorem is the following:

Theorem 1.3. *If we have a first order congruence of $\mathbb{P}^n$ such that the fundamental locus F satisfies the hypothesis of the preceding theorem, then F cannot be a complete intersection. If moreover F is smooth, then $n \leq 5$.*

Proof. In fact, by the preceding theorem $\deg(F) < (n-1)^2$, therefore if it were a complete intersection, it would be contained in a hypersurface V of degree less than $n-1$, and so every $(n-1)$ -secant line of F would be contained in V .

If F is smooth, since by [HS85] we know that Hartshorne's conjecture is true in codimension two up to degree $(n-1)(n+5)$, then $\dim(F) \leq 3$. $\square$

2. GENERAL EXAMPLES OF FIRST ORDER CONGRUENCES

We give now two examples of first order congruences of lines of $\mathbb{P}^n$. Actually, these examples give us all the congruences of Theorem 0.1 but case 4b.

2.1. Linear sections of $\mathbb{G}(1, n)$. First of all, we will analyse the congruences that come out from linear sections of the Grassmannian $\mathbb{G}(1, n)$, *i.e.* we will consider the so called, classically, *linear congruences*.

We recall that the Schubert cycle that corresponds to a hyperplane section of (the projective embedding of) the Grassmannian is σ_1 , so the following technical lemma gives us the formula for the general intersection of these special Schubert cycles:

Lemma 2.1. *If $\ell \leq n-1$ and we set $k := \lfloor \frac{\ell}{2} \rfloor$, the following formula holds:*

$$(4) \quad \sigma_1^\ell = \sum_{i=0}^k \left(\binom{\ell-1}{i} - \binom{\ell-1}{i-2} \right) \sigma_{\ell-i, i}$$

—with the convention that $\binom{\ell}{h} = 0$ if $h < 0$.

Proof. Let us prove the lemma by induction: for $\ell = 1$ it is obvious. Let us suppose it is true for $\ell-1$; then, by inductive hypothesis

$$\sigma_1^{\ell-1} = \sum_{i=0}^{k'} \left(\binom{\ell-2}{i} - \binom{\ell-2}{i-2} \right) \sigma_{\ell-1-i, i},$$

where $k' := \lfloor \frac{\ell-1}{2} \rfloor$. By Pieri's formula, we have

$$\sigma_{\ell-1-i, i} \cdot \sigma_1 = \begin{cases} \sigma_{\ell-i, i} & \text{if } \ell-1-i = i \\ \sigma_{\ell-i, i} + \sigma_{\ell-1-i, i+1} & \text{otherwise,} \end{cases}$$

i.e. if $\ell-1 \neq 2k'+1$ and $i \neq k'$, we obtain

$$\binom{\ell-2}{i} - \binom{\ell-2}{i-2} + \binom{\ell-2}{i-1} - \binom{\ell-2}{i-3} = \binom{\ell-1}{i} - \binom{\ell-1}{i-2}$$

while if $i = k' = \frac{\ell-2}{2}$,

$$\binom{2k'}{k'} - \binom{2k'}{k'-2} = \binom{2k'+1}{k'+1} - \binom{2k'+1}{k'-1},$$

which follows from the formula

$$(5) \quad \binom{\ell-1}{i} - \binom{\ell-1}{i-2} = \binom{\ell}{i} \cdot \frac{\ell-2i+1}{\ell-i+1}.$$

$\square$

Theorem 2.2. *If Λ is a congruence with sequence of degrees $(a_0, \dots, a_\nu)$ then B , as a subvariety of the Plücker embedding of the Grassmannian, has degree*

$$\deg(B) = \sum_{i=0}^{\nu} a_i \binom{n}{i} \cdot \frac{n-2i+1}{n-i+1}.$$

Proof. It is a corollary of formulae (4) and (5). $\square$

Corollary 2.3. *An $(n-1)$ -linear section B of the Grassmannian of lines of $\mathbb{P}^n$ generates a first order congruence Λ with sequence of degrees*

$$(\nu+1)\text{-deg}(\Lambda) = \left(1, \dots, \left(\binom{n-2}{i} - \binom{n-2}{i-2}\right), \dots, \left(\binom{n-2}{\nu} - \binom{n-2}{\nu-2}\right)\right);$$

in particular, as a subvariety of the Plücker embedding of the Grassmannian, this is a smooth congruence of degree

$$\deg(B) = \sum_{j=0}^{\nu} \left(\binom{n-2}{j} - \binom{n-2}{j-2} \right)^2 = \frac{1}{n-1} \binom{2n-2}{n}.$$

Obviously the degree of a linear congruence is the degree of the Grassmannian (which can be found in [Ful84], Example 14.7.11). This corollary gives us a first non-trivial example of a first order congruence. Some general results about fundamental varieties of these congruences are given in [BM01]; in particular, it is proven that the focal locus is the degeneracy locus F of a general morphism

$$(6) \quad \phi : \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)} \rightarrow \Omega_{\mathbb{P}^n}(2)$$

of (coherent) sheaves on $\mathbb{P}^n$ and that F is smooth if $\dim(F) \leq 3$.

The idea of the proof is the following: first we have the isomorphism:

$$(7) \quad H^0(\Omega_{\mathbb{P}^n}(2)) \cong (\wedge^2 V)^*,$$

where $V := H^0(\mathcal{O}_{\mathbb{P}^n}(1))^*$; from this we can interpret a global section of $\Omega_{\mathbb{P}^n}(2)$ as a bilinear alternating form on V , or as a skew-symmetric matrix of type $(n+1) \times (n+1)$ with entries in the base field. Then, the general morphism ϕ defined in (6) is assigned by giving $n-1$ general skew-symmetric matrices, $A_1, \dots, A_{n-1}$ and the corresponding degeneracy locus F in $\mathbb{P}^n$ is the set of points P such that

$$(8) \quad \sum_{i=1}^{n-1} \lambda_i A_i[P] = 0$$

for some $(\lambda_1, \dots, \lambda_{n-1}) \neq (0, \dots, 0)$, and $[P]$ denotes the column matrix of the coordinates of P . Finally, recalling that the Plücker embedding of the Grassmannian is:

$$\psi : \mathbb{G}(1, n) \hookrightarrow \mathbb{P}(\wedge^2 V)$$

we can conclude easily.

An improvement of a result of [BM01] is the following:

Proposition 2.4. *If F is the focal locus of a general linear congruence of $\mathbb{P}^n$, then*

1. *if n is even, equation (8) has at least one solution and F is a rational variety;*
2. *if n is odd, the vanishing of the Pfaffian of the matrix of (8) defines a hypersurface Z of degree $(n+1)/2$ contained in a $\mathbb{P}^{n-2}$ in which $\lambda_1, \dots, \lambda_{n-1}$ are the coordinates. Furthermore, if ϕ is general, for a fixed point $[\lambda] \in Z$, we find a line of solutions of equation (8) in F , so that F is a scroll over (an open set of) Z .*

Besides,

$$\deg(F) = \frac{n^2 - 3n + 4}{2}.$$

Proof. If we dualise the Eagon-Northcott complex applied to (6), we have the following exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(1-n) \rightarrow \mathcal{T}_{\mathbb{P}^n}(-2) \xrightarrow{t\phi} \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)} \rightarrow \omega_F^\circ(2) \rightarrow 0,$$

where ω_F° is the dualising sheaf of F . Hunting in the sequence, we get

$$H^0(\mathbb{P}^n, \omega_F^\circ(2)) \cong H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}),$$

and so the complete linear system $|K_F + 2H|$ —where K_F and H are the canonical and hyperplane divisors, respectively—is base point free, from which we obtain that the map associated to it

$$\varphi_{|K_F + 2H|} : F \rightarrow \mathbb{P}^{n-2}$$

is well defined. The fibre of $\varphi_{|K_F + 2H|}$ —if we fix projective coordinates $\lambda_1, \dots, \lambda_{n-1}$ on $\mathbb{P}^{n-2}$ and $x_0 \dots x_n$ on $\mathbb{P}^n$ —is given, in view of (8), by the solutions of the following homogeneous linear system of $n+1$ equations in the $n+1$ indeterminates: $x_0 \dots x_n$:

$$(9) \quad \sum_{\substack{i=1, \dots, n-1 \\ k=0, \dots, n}} \lambda_i (A_i)_{j,k} x_k = 0, \quad j = 0, \dots, n,$$

where $\lambda_1, \dots, \lambda_{n-1}$ are fixed. Clearly the $(n+1) \times (n+1)$ -matrix $A := \sum \lambda_i (A_i)$ associated to this system is antisymmetric; therefore, if n is even, its determinant is zero and has only one or infinitely many (projective) solutions; so, for dimensional reasons, we get that $\varphi_{|K_F + 2H|}$ is birational and F is rational.

If instead n is odd, $\det A$ is, in general, not zero, and so its Pfaffian defines a hypersurface Z of degree $(n+1)/2$ and the general fibre of $\varphi_{|K_F + 2H|}$ is a $\mathbb{P}^1$. We observe that the Eagon-Northcott gives in fact a locally free resolution of the ideal sheaf of F :

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}(1-n) \xrightarrow{\phi(1-n)} \Omega_{\mathbb{P}^n}(3-n) \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \mathcal{O}_F \rightarrow 0.$$

From this we get the Hilbert polynomial of F , and in particular, after some computations, the degree. $\square$

Remark. In low dimension and with a general section, we have that (see [BM01] for details) if $n = 3$, F is the union of two skew lines, if $n = 4$, F is a smooth projected Veronese surface and if $n = 5$, F is a (rational) threefold of degree seven, which is a scroll over a cubic surface in $\mathbb{P}^3$. It is also known as Palatini scroll (see [Ott92]).

Remark. Since a linear congruence B is given by a linear section of the Grassmannian $\mathbb{G}(1, n)$, a resolution of its ideal sheaf is given by the Koszul complex

$$\begin{aligned} 0 \rightarrow \mathcal{O}_{\mathbb{G}(1,n)}(-n+1) \rightarrow (\wedge^2((\mathcal{O}_{\mathbb{G}(1,n)}(-1))^{\oplus(n-1)})) \rightarrow \dots \\ \dots \rightarrow ((\mathcal{O}_{\mathbb{G}(1,n)}(-1))^{\oplus(n-1)}) \rightarrow \mathcal{O}_{\mathbb{G}(1,n)} \rightarrow \mathcal{O}_B \rightarrow 0. \end{aligned}$$

Instead, the minimal free resolution of the Plücker embedding of B can be obtained by the minimal free resolution of $\mathbb{G}(1, n)$ —which can be found in [JPW81]—via the mapping cone. Since the computations are rather complicate, we will not give them.

2.2. Matrices of type $(n-1) \times n$ with linear entries. Let us consider a general morphism $\phi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}, \mathcal{O}_{\mathbb{P}^n}^{\oplus n}(1))$, whose minors vanish in the expected codimension two. In this case, $F := V(\phi)$ —the degeneracy locus of ϕ —is a locally Cohen-Macaulay subscheme, the Eagon-Northcott complex is exact (see [BE75]) and gives a free resolution of our ideal sheaf:

$$(10) \quad 0 \rightarrow \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}(-n) \xrightarrow{\phi(-n)} \mathcal{O}_{\mathbb{P}^n}^{\oplus n}(1-n) \rightarrow \mathcal{O}_{\mathbb{P}^n} \rightarrow \mathcal{O}_F \rightarrow 0.$$

Then—for example—from the Hilbert polynomial we get

$$(11) \quad \deg(F) = \binom{n}{2}$$

$$(12) \quad \pi(F) = 1 + \frac{2n-7}{3} \binom{n}{2}$$

where $\pi(F)$ is the sectional genus of F . It is easy to prove that:

Proposition 2.5. *F is rational, and if $n \leq 5$ it is smooth. Besides, the adjunction map $\varphi_{|K_F+H|}$ exhibits F as the blow-up of $\mathbb{P}^{n-2}$ in a scheme Z of degree $\binom{n+1}{2}$ and sectional genus $\frac{n}{6}(2n-5)(n+1)-1$. In particular, if $n=4$, F is a rational sextic which is the blow-up of the plane in 10 points, i.e. a Bordiga surface.*

Proof. The smoothness is a consequence of Bertini type theorems. For proving that F is rational, we can apply the standard argument used in the proof of Proposition 2.4: we consider the dual of the Eagon-Northcott complex applied to ϕ

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-n) \rightarrow \mathcal{O}_{\mathbb{P}^n}^{\oplus n}(-1) \xrightarrow{t\phi} \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)} \rightarrow \omega_F^\circ(1) \rightarrow 0,$$

where ω_F° is the dualising sheaf of F . Chasing in the sequence, we get

$$H^0(\mathbb{P}^n, \omega_F^\circ(1)) \cong H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}^{\oplus(n-1)}),$$

and the complete linear system $|K_F + H|$ —where K_F and H are the canonical and hyperplane divisors, respectively—is base point free, from which we obtain that the map associated to it

$$\varphi_{|K_F+H|} : F \rightarrow \mathbb{P}^{n-2}$$

is well defined. The fibre of $\varphi_{|K_F+H|}$ —if we fix projective coordinates $y_0, \dots, y_{n-2}$ on $\mathbb{P}^{n-2}$ and $x_0 \dots x_n$ on $\mathbb{P}^n$ —is given by the solutions of the following homogeneous linear system of n equations in the $n+1$ indeterminates $x_0 \dots x_n$:

$$(13) \quad \sum_{i=0}^n \Phi_{j,i}(y_0 : \dots : y_{n-2})x_i = 0, \quad j = 1, \dots, n,$$

where $y_0, \dots, y_{n-2}$ are fixed and $\Phi_{j,i}(y_0 : \dots : y_{n-2})$ are linear forms on $\mathbb{P}^{n-2}$. Clearly the $(n+1) \times n$ -matrix associated to this system has maximal rank, since the forms $\Phi_{j,i}(y_0 : \dots : y_{n-2})$ are linearly independent if ϕ is general, and then it has only one (projective) solution; so, $\varphi_{|K_F+H|}$ is birational and F is rational. Besides, $\varphi_{|K_F+H|}$ has (at least) one dimensional fibres on the degeneracy locus Z of the map $\Phi : \mathcal{O}_{\mathbb{P}^{n-2}}^{\oplus n} \rightarrow \mathcal{O}_{\mathbb{P}^{n-2}}^{\oplus(n+1)}(1)$, i.e. $\varphi_{|K_F+H|}$ gives F as the blow-up of $\mathbb{P}^{n-2}$ in Z . As usual, Eagon-Northcott gives a free resolution of Z :

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^{n-2}}^{\oplus n}(-n-1) \rightarrow \mathcal{O}_{\mathbb{P}^{n-2}}^{\oplus(n+1)}(-n) \rightarrow \mathcal{O}_{\mathbb{P}^{n-2}} \rightarrow \mathcal{O}_Z \rightarrow 0,$$

from which we obtain the degree and the sectional genus of Z . $\square$

With this, we prove that

Theorem 2.6. *The $(n-1)$ -secant lines of the variety F defined as above form a first order congruence of lines B of $\mathbb{P}^n$. The congruence B is smooth for general ϕ .*

Proof. Let

$$A := \begin{pmatrix} a_{11} & \cdots & a_{1(n-1)} \\ \vdots & & \\ a_{n1} & & a_{n(n-1)} \end{pmatrix}$$

be the matrix of linear entries on $\mathbb{P}^n$ associated to the map ϕ . Moreover, let us denote by A_i , $i = 1, \dots, n$ the rows of A and by a_i the minor of order $n-1$ of A obtained cutting out the row A_i . We want to prove that any $(n-1)$ -secant line of F is given by the vanishing of the entries of $A_\lambda := \sum_{i=1}^n \ell_{\lambda,i} A_i$, i.e. by a linear combination of the rows of A : $\ell_\lambda := V(\sum_{i=1}^n \ell_{\lambda,i} a_{ij})$, $j = 1, \dots, n-1$. In fact, ℓ_λ is an $(n-1)$ -secant line of F : after changing coordinates, we can assume that ℓ_λ is defined by the zeros of the last row. The intersection $V(a_n) \cap \ell_\lambda$ is given by $n-1$ points $P_1, \dots, P_{n-1}$ and by definition, the matrix A drops rank at each of the P_i 's.

Vice versa, if ℓ is an $(n-1)$ -secant line, to prove that its equations are a linear combination of rows it is sufficient to show that ℓ is contained in at least one (linear combination) of the hypersurfaces $V(a_i)$. If this were not so, we would have that $V(a_i) \cap \ell = F \cap \ell = \{P_1, \dots, P_{n-1}\}$, $\forall i$, since $V(a_i) \cap \ell \supset F \cap \ell$ and both sets contain $n-1$ points. But if we take a pencil of these hypersurfaces, we can find a linear combination of these containing ℓ .

By this, it is easy to see that B is a first order congruence: if $P \in \mathbb{P}^n$ is a general point, we have to solve $A(P) \cdot {}^t \lambda$ with $\lambda \in \mathbb{P}^{n-1}$, *i.e.* a homogeneous linear system of $n-1$ equations in n indeterminates.

Finally, for proving that B is smooth, from the above geometric description of the $(n-1)$ -secant lines as linear combination of the rows of A , we see that the map ϕ gives rise to a map

$$\varphi : \mathcal{O}_{\mathbb{G}(1,n)}^{\oplus n} \rightarrow (\mathcal{S}^*)^{\oplus (n-1)}.$$

—where φ is obtained by considering the dual of ϕ twisted by one and then pulled back to the incidence variety and finally pushed it forward to $\mathbb{G}(1,n)$ —and B is precisely the degeneracy locus of φ . Then we conclude as usual by Bertini type theorems. $\square$

Remark. If we apply the Eagon-Northcott complex to φ , we obtain a resolution of B :

$$(14) \quad 0 \rightarrow \mathcal{O}_{\mathbb{G}(1,n)}^{\oplus \binom{n}{2}} \rightarrow (\mathcal{S}^*)^{\oplus (n-1) \binom{n}{3}} \rightarrow \\ \rightarrow (\wedge^2((\mathcal{S}^*)^{\oplus (n-1)}))^{\oplus \binom{n}{4}} \rightarrow \dots \\ \dots \rightarrow \wedge^{n-2}((\mathcal{S}^*)^{\oplus (n-1)}) \rightarrow \mathcal{O}_{\mathbb{G}(1,n)}(n-1) \rightarrow \mathcal{O}_B(n-1) \rightarrow 0.$$

3. TWO MULTIPLE POINT FORMULAE

We will prove now the quadruple point formula for a smooth threefold X of $\mathbb{P}^5$ and the formula which gives the number of 4-secant lines to a surface S of $\mathbb{P}^4$ passing through a general point $P \in S$.

In this and in the next section, we will denote by S a general hyperplane section of the threefold X , by H and K the hyperplane and the canonical divisors of X . $d := \deg(X) = H^3$ is the degree of X , $\pi = \frac{1}{2}H^2(K + 2H) + 1$ its sectional genus, while $\chi(\mathcal{O}_X)$ and $\chi(\mathcal{O}_S)$ are, respectively, the Euler-Poincaré characteristic of X and S . We recall that for a smooth threefold of $\mathbb{P}^5$ two double point formulae hold—*i.e.* one for X and one S —which can be written as

$$K^3 = -5d^2 + d(2\pi + 25) + 24(\pi - 1) - 36\chi(\mathcal{O}_X) - 24\chi(\mathcal{O}_S), \\ H \cdot K^2 = \frac{1}{2}d(d+1) - 9(\pi - 1) + 6\chi(\mathcal{O}_X),$$

so the basic invariants of X are $d, \pi, \chi(\mathcal{O}_X)$ and $\chi(\mathcal{O}_S)$ (see for example [DP95]).

We start with the quadruple point formula. We refer to [Kle81] for the definitions and results used in the proof.

Proposition 3.1. *Let X be a smooth threefold of $\mathbb{P}^5$ and $P \in (\mathbb{P}^5 \setminus X)$ is a point through which there is a finite number of 4-secant lines of X . If we set $q(X) := \text{length}(\Psi \cap W_P)$ where $\Psi, W_P \subset \mathbb{G}(1,n) \subset \mathbb{P}^{14}$ are, respectively, the family of 4-secant lines of X and the Schubert variety of the lines through P (embedded via the Plücker embedding), then the following formula holds:*

$$(15) \quad q(X) = \frac{1}{24}d^4 - \frac{1}{4}d^3 + \frac{1}{2}d^2\left(\frac{11}{12} - \pi\right) + d\left(\frac{5}{2}\pi + 2\chi(\mathcal{O}_S) - \frac{9}{4}\right) + \\ + \frac{1}{2}\pi^2 - \frac{7}{2}\pi + 6\chi(\mathcal{O}_X) - 9\chi(\mathcal{O}_S) + 3,$$

In particular, for the general point of $\mathbb{P}^5$, there pass $q(X)$ 4-secant lines, i.e. Ψ —if it is not empty—is a (possibly reducible) congruence of order $a_0 = q(X)$.

Proof. Let us consider the projection $\pi_P : X \rightarrow \mathbb{P}^4$ from a general point P of $\mathbb{P}^5$ to a hyperplane. π_P is—as observed in [Kle81]—4-generic (see [Kle81] for the definition). Therefore, we can apply Kleiman’s quadruple point formula of [Kle81], and from this we obtain formula (15) in the case of a general point.

If P is not a general point, the conclusion follows by degree reasons. $\square$

Next, we pass to prove the other formula.

Proposition 3.2. *Let $S \subset \mathbb{P}^4$ be a smooth non-scalar surface of degree d , sectional genus π , hyperplane and canonical divisors H and K , respectively. Then the number of 4-secant lines h of S passing through a general point $P \in S$ is given by the formula*

$$(16) \quad h = \frac{1}{6}d^3 - \frac{3}{2}d^2 + d\left(\frac{16}{3} - \pi\right) + 4\pi + 2\chi(\mathcal{O}_S) - 10.$$

Proof. h is actually equal to the number of triple points of the image of S under the projection from P to a hyperplane. Therefore h can be obtained from the triple point formula for a map f from a smooth surface to $\mathbb{P}^3$: we simply blow-up S in P and then we compose the map which defines the blow-up $g : \text{Bl}_P(S) \rightarrow S$ with the projection, i.e. $f := \pi_P \circ g$. The triple point formula can be found in [Le 87], and it is, in our situation

$$(17) \quad h = \frac{1}{6}(\tilde{d}(\tilde{d}^2 - 12\tilde{d} + 44) + 4\tilde{K}^2 - 2\tilde{c}_2 - 3\tilde{H}\tilde{K}(\tilde{d} - 8)),$$

where $\tilde{H} = g^*H - E$ is the strict transform of H — E is the exceptional divisor of the blow-up—, $\tilde{K} = g^*K + E$ the canonical divisor of $\text{Bl}_P(S)$, $\tilde{d} = \tilde{H}^2$ and $\tilde{c}_2$ its topological Euler-Poincaré characteristic. Clearly, we have that $\tilde{d} = d - 1$, $\tilde{c}_2 = c_2 + 1$, where c_2 is the topological Euler-Poincaré characteristic of S , $\tilde{K}^2 = K^2 - 1$ since S is not a scroll, and $\tilde{H}\tilde{K} = 2\pi - d - 1$. Then, if we express the invariants of S in terms of the basic invariants $(d, \pi, \chi(\mathcal{O}_S))$, we get formula (16). $\square$

Remark. We obtained the formulae with the help of S. Katz and S. A. Strømme’s Maple package “Schubert”.

4. CONGRUENCES OF $\mathbb{P}^n$

In this section we study the irreducible first order congruences B that are given by the families of the 4-secant lines of smooth threefolds $X \subset \mathbb{P}^5$ (with the notations for its invariants given in Section 3), proving Theorem 0.1. We need the following preliminary result:

Lemma 4.1. *If $S(= X \cap H)$ is not a scroll, the following formula holds:*

$$(18) \quad 0 = \frac{1}{8}d^4 - \frac{23}{12}d^3 - d^2\left(\pi - \frac{83}{8}\right) - d\left(\frac{355}{12} - 11\pi - 2\chi(\mathcal{O}_S)\right) + \frac{1}{2}\pi^2 - \frac{57}{2}\pi - 17\chi(\mathcal{O}_S) + 53.$$

Proof. Formula (18) is formula (1), which is, in our situation,

$$(19) \quad 4h = 1 + a_1$$

(with the notations of the proof of Theorem 1.2). Now, h is the algebraic multiplicity of the fundamental locus on the variety V_Π of the lines of B that meet a general 3-plane Π , i.e. if we fix $P \in X$, there are h lines of B though it that meet Π also. The hyperplane $\overline{P\Pi}$ intersects X in a smooth surface S , and the h lines are exactly the ones that are 4-secants to S . So h is given by formula (16).

By definition, a_1 is the number of lines of B contained in a hyperplane H and that meet a line $\ell \subset H$, *i.e.* it is the degree of the hypersurface of the 4-secant lines of S . But this formula can be easily deduced from [Le 90] and it is

$$(20) \quad a_1 = \frac{1}{8}d^4 - \frac{5}{4}d^3 + d^2\left(\frac{35}{8} - \pi\right) + d\left(+7\pi + 2\chi(\mathcal{O}_S) - \frac{33}{4}\right) + \frac{1}{2}\pi^2 - \frac{25}{2}\pi - 9\chi(\mathcal{O}_S) + 12.$$

Substituting formulae (16) and (20) in (19), we get formula (18). $\square$

Remark. During the proof of the preceding proposition, we have calculated the first multidegree a_1 for the family (possibly reducible) of the 4-secant lines of a smooth threefold X of $\mathbb{P}^5$. a_2 is instead the number of 4-secant lines contained in a 3-dimensional linear space G , *i.e.* the number of 4-secant lines of the smooth curve $G \cap X$, and this formula is in [Le 82]:

$$(21) \quad a_2 = \frac{1}{12}d^4 - d^3 + \frac{53}{12}d^2 - \frac{17}{2}d + 6 - \frac{1}{2}\pi d^2 + \frac{7}{2}d\pi - \frac{13}{2}\pi + \frac{1}{2}\pi^2.$$

Lemma 4.2. *The family B of the 4-secant lines of the threefold X of case (4b) of Theorem 0.1 is irreducible.*

Proof. First of all, recall that $X = \cup_{\ell \in Z} \ell$, where $Z := \mathbb{G}(1, 5) \cap \mathbb{P}^8 (\subset \mathbb{P}^{14})$: see Proposition 3.2 of [Cha89].

Applying Formula (15), we get that B is a first order congruence. If B were reducible, we could write $B = B_0 \cup B_1$, where B_1 is the irreducible component of B of order 1 and B_0 is the union of the other components (which have order zero); besides, we denote by $a_j(i)$ the j -th multidegree of B_i . Now, Formula (19) becomes $4h = 1 + a_1(1)$, but in Formula (20) we have to substitute a_1 with $a_1(0) + a_1(1)$, and therefore $a_1(0) = 0$. So, the only possibility is $[B_0] = a_2(0)\sigma_{2,2}$, *i.e.* X contains at least a surface of a $\mathbb{P}^3$ of degree at least 4.

If $Y \subset \Pi \cong \mathbb{P}^3$ were one of these surfaces, it is enough to prove that Y is not a scroll. In fact, once we have proven this, we can see that Y would be birationally equivalent to Z , since through almost every point of Y there is a line of X not contained in Π . But Z is the intersection of $\mathbb{G}(1, 5)$ with six general hyperplanes of $\mathbb{P}^{14}$, while in $\mathbb{P}^{14*}$ the set of hyperplanes consisting of the special Schubert cycles of lines meeting a 3-plane has codimension six (it is in fact $\mathbb{G}(3, 5)$) and hence disjoint with the span of six general elements.

If Y were a scroll, then its corresponding curve C in X would be contained in a fixed $\mathbb{G}(1, \Pi) \subset \mathbb{G}(1, 5)$. But both X and $\mathbb{G}(1, \Pi)$ are linear sections of $\mathbb{G}(1, 5)$ and therefore C could be (at most) a conic, and so Y could only be a quadric. $\square$

Remark. In the previous proof, the fact that we can exclude the non-scrollar case, it has been suggested to us by the referee. Our proof was longer and based on the fact that the sectional genus of Y would be the same of Z .

Proof of Theorem 0.1. By Theorem 1.3, it is enough to consider the cases of $\mathbb{P}^n$ with $n \leq 5$.

By Theorem 1.2 (and by Castelnuovo's bound) we get that in $\mathbb{P}^3$ the only congruence is the one given by the secant lines of the twisted cubic.

In $\mathbb{P}^4$, again by Theorem 1.2, we obtain that the surfaces X we are looking for have to satisfy $4 \leq \deg(X) \leq 8$, and since the smooth surfaces are classified up to degree ten, see [DP95], if we apply the triple point formula (see [Le 87], or formula (17), erasing the tildes) to them, we get only the cases of the theorem.

Passing to the next case, since the smooth threefolds of $\mathbb{P}^5$ are classified up to degree 12 (see [BSS95]), we can check which of them have an apparent quadruple point, and it turns out that they are the ones of the list of the theorem. The three

congruences are indeed irreducible, case (4b) by Lemma 4.2, the other two from their description as degeneracy loci.

Next, from Theorem 1.2 we have to exclude the cases of degrees 13, 14 and 15. To do this, we calculate the possible invariants of these threefolds (see for example [BSS95]) and then we request that they have to satisfy $q(X) = 1$ in formula (15), and equation (18) applied to the general hyperplane section S of X : since the smooth, non-degenerate scroll surfaces of $\mathbb{P}^4$ have degree at most 5, see [Lan80], we can apply this last formula without problems. It turns out that there cannot exist threefolds with these conditions.

Finally we can calculate the multidegree for the three congruences of $\mathbb{P}^5$ (the cases of $\mathbb{P}^3$ and $\mathbb{P}^4$ can be easily deduced from Section 2); a_1 is from formula (20) and a_2 is given by formula (21). $\square$

Remark. We performed the calculations in the last proof with the help of a simple program in Maple.

REFERENCES

- [BE75] D. A. Buchsbaum and D. Eisenbud, *Generic free resolutions and a family of generically perfect ideals*, Adv. Math. **18** (1975), 245–301.
- [BM01] D. Bazan and E. Mezzetti, *On the construction of some Buchsbaum varieties and the Hilbert scheme of elliptic scrolls in $\mathbb{P}^5$* , Geom. Dedicata **86** (2001), no. 1–3, 191–204.
- [BSS95] M. Beltrametti, M. Schneider, and A. Sommese, *Some special properties of the adjunction theory for 3-folds in $\mathbb{P}^5$* , Mem. Am. Math. Soc. **554** (1995), 63 p.
- [Cha89] M. Chang, *Classification of Buchsbaum subvarieties of codimension 2 in projective space*, J. Reine Angew. Math. **401** (1989), 101–112.
- [CS89] C. Ciliberto and E. Sernesi, *Families of varieties and the Hilbert scheme*, Riemann surfaces (proceedings of the 1987 ICTP college on Riemann surfaces) (M. Cornalba et al., eds.), ICTP, World Scientific Press, 1989, pp. 428–499.
- [De 99] P. De Poi, *On first order congruences of lines*, Ph.D. thesis, SISSA-ISAS, Trieste, Italy, October 1999, <http://www.dm.unipi.it/~depoi/pubblicazioni/focchi.ps>.
- [De 01] P. De Poi, *On first order congruences of lines of $\mathbb{P}^4$ with a fundamental curve*, Man. Math. **106** (2001), 101–116.
- [DP95] W. Decker and S. Popescu, *On surfaces in $\mathbb{P}^4$ and 3-folds in $\mathbb{P}^5$* , Vector bundles in algebraic geometry. Proceedings of the 1993 Durham symposium, Durham, UK. (N. J. Hitchin et al., eds.), Lond. Math. Soc. Lect. Note Ser., no. 208, Cambridge University Press, 1995, pp. 69–100.
- [Ede94] G. Edelmann, *3-folds in $\mathbb{P}^5$ of degree 12*, Man. Math. **82** (1994), no. 3–4, 393–406.
- [Eis95] D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*, Graduate Texts in Mathematics, no. 150, Springer-Verlag, Berlin, 1995.
- [Ful84] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag, 1984.
- [GH78] P. Griffiths and J. Harris, *Principles of algebraic geometry*, John Wiley & Sons, 1978.
- [Har77] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, no. 52, Springer-Verlag, New York - Heidelberg - Berlin, 1977.
- [HS85] A. Holme and M. Schneider, *A computer aided approach to codimension 2 subvarieties of $\mathbb{P}^n$, $n \geq 6$* , J. Reine Angew. Math. **357** (1985), 205–220.
- [JPW81] T. Jozefiak, P. Pragacz, and J. Weyman, *Resolutions of determinantal varieties and tensor complexes associated with symmetric and antisymmetric matrices*, Tableaux de Young et foncteurs de Schur en algèbre et géométrie. Conference internationale, Torun, Pologne, 27 aout - 3 septembre 1980, Astérisque, no. 87–88, Société Mathématique de France, 1981, pp. 109–189.
- [Kle77] S. L. Kleiman, *The enumerative theory of singularities*, Real and complex singularities, Oslo 1976 (P. Holm, ed.), Sijthoff & Noordhoff International Publishers, 1977, pp. 297–396.
- [Kle81] S. L. Kleiman, *Multiple-point formulas. I: Iteration*, Acta Math. **147** (1981), 13–49.
- [Kwa01] S. Kwak, *Smooth threefolds in $\mathbb{P}^5$ without apparent triple or quadruple points and a quadruple-point formula*, Math. Ann. **320** (2001), no. 4, 649–664.
- [Lan80] A. Lanteri, *On the existence of scrolls in $\mathbb{P}^4$* , Atti Accad. Naz. Lincei, VIII. Ser., Rend., Cl. Sci. Fis. Mat. Nat. **69** (1980), 223–227.

- [Le 82] P. Le Barz, *Formules multisécantes pour les courbes gauches quelconques*, Enumerative geometry and classical algebraic geometry, Nice 1981 (P. Le Barz and Y. Hervier, eds.), Progress in Mathematics, no. 24, Birkhäuser, 1982, pp. 165–197.
- [Le 87] P. Le Barz, *Formules pour les trisécantes des surfaces algébriques*, L'Enseignement Mathématique **33** (1987), 1–66.
- [Le 90] P. Le Barz, *Quelques formules multisécantes pour les surfaces*, Enumerative geometry (Sitges, 1987) (Berlin) (S. Xambó-Descamps, ed.), Lecture Notes in Mathematics, no. 1436, Springer-Verlag, Berlin, 1990, pp. 151–188.
- [Ott92] G. Ottaviani, *On 3-folds in $\mathbb{P}^5$ which are scrolls*, Ann. Scuola Norm. Sup. Pisa, cl. Sci. **19** (1992), 451–471.